

The epigeic spider communities of lowland forests in the surroundings of the Danube River on the territory of Slovakia and their usage for biota monitoring

PETER GAJDOŠ

Institute of Landscape Ecology, Slovak Academy of Sciences,
Akademická 2, 949 01 Nitra, Slovak Republic

Gabčíkovo-Nagymaros water dam, inundation forests, Danube River, spider communities, monitoring

Abstract. Research on spiders was carried out on seven selected model areas of inundation forests along the Danube River during 1991–1993. Altogether 2240 individual spiders were caught, belonging to 104 species. In the wet soft wood inundation forests, *Pirata hygrophilus* was eudominant ($D > 40\%$). On the drier sites *Ozyptila praticola* was eudominant ($D > 29\%$). The proportion in which these two species are present in soft wood inundation forests is a very good marker for moisture evaluation of the model areas. Changes in the spider communities were observed during the time of research in consequences of the Gabčíkovo-Nagymaros dam construction and functioning.

INTRODUCTION

The Danube River on the boundary of Slovakia and Hungary, creates an inland delta with a motley of biotopes and a high biodiversity. Due to the evident influence of human activities upon the surrounding landscape, such as the Gabčíkovo-Nagymaros dam construction and functioning, the need arose for a long term biota monitoring and for regular evaluation and prognosis of the situation in that area. The research of the above mentioned territory started in 1991, two years before the dam began to function, on selected model areas along the Danube River.

The first information on spiders from the model areas were published by Böckh (1857, 1862a,b) and Ortway (1902), and come from the surroundings of Bratislava. Some sporadic data from the last century were given by Chyzer and Kulczyński (1891, 1894, 1897, 1899). Gajdoš (1987) investigated the spiders in the State Nature Reservation Kopáč which is situated very close to the Danube River. Krumpálová (1994) and Gajdoš (1994b) were dealing with selected areas along the Danube River. A summary of the present time knowledge about the spiders of the Podunajská Nižina lowlands on the Slovakia territory was published by Gajdoš et al. (1992).

Data on spider communities of inundation forests along the Danube River from Austria and Germany were published by Thaler et al. (1984), Thaler and Steiner (1989) and Bauchhenss (1991), respectively.

MATERIAL AND METHODS

The research of the spiders was carried out on six selected model areas during 1991 and 1992, which were in 1993 supplemented with one additional locality (Čičov) (Fig. 1). The spiders were collected by the square method (square 25 x 25 cm) from 16 different squares, altogether 1 m². The samples were collected about once a month, in the period from March till November from each locality. The spiders were taken out of the samples by a xerolector (Mocarsky type).

The similarity of the communities was evaluated on the basis of the Wishart index. Cluster analysis was performed by program NCLAS2 (Podanini, 1980) with the use of the UPGMA method. The spiders collected are listed (Table 1) in the order of Platník (1993). Average yearly soil moisture was calculated from two-weekly measurements, to 10 cm of soil depth (Janík 1991–1994).

OVERVIEW OF THE MODEL AREAS

Bo—Bodíky

Inundation forest (*Saliceto-Populetum*) in the surroundings of the village Bodíky, age of trees 80 years (river kilometer 1830, 115 m altitude above sea level).




Fig. 1. Position of the model areas in surroundings of the Danube river.

Ga—Gabčíkovo-Istragov

Inundation forest (*Saliceto-Populetum*) in the surroundings of the town Gabčíkovo, age of trees 80 years (river kilometer 1817, 113 m altitude).

To—Trstená na Ostrove-Kráľovská Lúka

Inundation forest (*Saliceto-Populetum*) in the surroundings of the village Trstená na Ostrove, age of trees 90 years (river kilometer 1826, 117 m altitude).

Do—Dobrohošť

Inundation forest (*Saliceto-Populetum*) in the surroundings of the village Dobrohošť, age of trees 60 years (river kilometer 1840, 120 m altitude).

Kl—Klúčovec

Inundation forest (*Saliceto-Populetum*) in the surroundings of the village Klúčovec, age of trees 100 years (river kilometer 1804.5, 116 m altitude).

Či—Čičov

Inundation forest (*Saliceto-Populetum*) in the surroundings of the village Čičov, age of trees 100 years (river kilometer 1799–1801, 116 m altitude).

VI—Bratislava-Vlčie hrdlo

Hard wood inundation forest (*Ulmeto-Fraxinetum*) remotes from the main river-bed of the Danube River, age of trees 80 years (river kilometer 1864, 132 m altitude).

RESULTS AND DISCUSSION

On the searched territory, 136 samples were collected during the period 1991–1993. Altogether 2240 individual spiders were caught, belonging to 104 species. In the wet soft wood inundation forests *Pirata hygrophilus* was eudominant ($D > 40\%$), while on the drier sites the *Ozyptila praticola* was dominant ($D > 29\%$). *Pirata hygrophilus* according to Martin (1991), is considered to be hygrophilous or hemihygrophilous. The ecological requirements for *Ozyptila praticola* are rather hemihygrophil to hemixerophil by Martin (1991). In the hard wood inundation forests the following species were eudominant: *Enoplognatha thoracica* (D_{91-93} from 14.5 to 16.7%), *Zelotes apricorum* ($D_{92} = 11.7\%$, $D_{93} = 19.4\%$), and partly in 1991 also *Ozyptila praticola* ($D_{91} = 14.5\%$). From faunistic point of view, the captures of rare and threatened species occurring on the Red list of spiders of Slovakia (Gajdoš & Svatoň, 1993), such as *Ero aphana*, *Anelosimus vittatus*, *Entelecara erythropus*, *Gongylidiellum murcidum*, *G. vivum*, *Lessertinella kulczynskii*, *Walckenaeria obtusa*, *Hahnia helveola*, *Anyphaena furva* and *Zelotes gracilis* are the most important (Table 1). The average annual abundance of spiders on 1 m² ranges from 7.2 (VI in 1993) to 28.7 specimens (To in 1992).

Gabčíkovo-Istragov

Belongs to the wetter site with a high abundance of hygrophilous species *Pirata hygrophilus* (D from 42.4 to 52.9%). Lisický et al. (1988) forecasted the decrease of the level of ground water in the crucial period 1990–1991. The communities at the time of the research showed stability (Shannon's index $H = 1.96\text{--}2.10$) with a slight increase in the number and densities of species. The rare species *Gongylidiellum murcidum* was present here.

Trstená na Ostrove-Kráľovská Lúka

Rather wet locality with a predominance of the hygrophilous species *Pirata hygrophilus* ($D = 42.0\text{--}55.2\%$). During the three years of research the communities showed stability ($H = 1.89\text{--}2.16$), which is in conformity with the prognosis of Lisický et al. (1988), who did not expect any changes due to the remoteness of the locality from the old river-bed. Species richness was a little bit lower as compared to the other localities (31 species). From the threatened species of the fauna of Slovakia, the following species were found: *Anelosimus vittatus* and *Gongylidiellum vivum*.

Dobrohošť

The spider communities showed changes in their composition during each of the three years. A considerable decrease was observed in the dominance of the hygrophilous species *Pirata hygrophilus* and on the other hand, an increase of the hemihygro-hemixerophilous species *Ozyptila praticola*. An increase of 50% in species richness was observed in 1993 in comparison with 1991. The other species which were found belong also rather to hemihygrophilous or hemixerophilous ones. It indicates a desiccation of the soil horizons and corresponds with the decrease of the soil moisture (Table 2).

Ključovec

After the Gabčíkovo-Nagymaros dam started to function on October 1992, the spider communities changed in favour of more hygrophilous species (considerable increase in the abundance of the species *Pirata hygrophilus* and decrease of the species *Ozyptila praticola* in 1993). Of faunistic interest are the captures of *Walckenaeria obtusa* and *Hahnina helveola*.

Bratislava-Vlčie hrdlo

This is the only model area of hard wood inundation forest which is reflected in the composition of the spider communities as compared to the soft wood inundation forests. There is a predominance of hemixerophilous species and the species *Pirata hygrophilus* is completely absent. The locality showed a relative stability during the period of investigation. From the 37 species three species are listed in the Red list of spiders of Slovakia: *Ero aphana*, *Anyphaena furva*, *Zelotes gracilis*.

Čičov

Locality with predominance of species *Ozyptila praticola* ($D = 38.46\%$) showed a rather dry character of the habitat. Average annual abundance of spiders on 1 m^2 was very low (10.8 specimens). Totally, 21 species of spiders were found.

COMPARISON OF EPIGEIC SPIDER COMMUNITIES

Based upon the similarity of the communities at the value of Wishart's index cca 23%, the yearly evaluated spider communities from each model area were divided into three large clusters by use of the UPGMA method (Fig. 2).

Cluster A represents the communities of more wet soft wood inundation forests (**Bo₉₁, Ga₉₂, Ga₉₁, To₉₂, To₉₃, Ga₉₃, To₉₁, Kl₉₃**) with eudominance of the species *Pirata hygrophilus*. The spider community of Klučovec (subcluster Ad) has an intermediate position in this cluster in 1993, when *Pirata hygrophilus* ($D = 26.3\%$) and *Ozyptila praticola* ($D = 20.9\%$) are equally numerous and in this way the community formed the link to the dryer areas of cluster B. Subclusters Aa, Ab and Ac show a presence of *Pirata hygrophilus* higher than 40%, at which level there is also a significant presence of *Ozyptila praticola* ($D > 16\%$) in the subcluster Aa, and of eudominant *Porrhomma*




Fig. 2. Hierarchical classification of epigeic spider communities in the model areas according to the similarity of abundance (Wishart's index (UPGMA)).

convexum in the community Ac ($D = 18.5\%$). The subcluster Ab represents the localities with the highest number of hygrophilous species, *Pirata hygrophilus* being the most frequent species ($D > 49\%$).

Cluster B represents the drier localities of soft wood inundation forests, where *Ozyptila praticola* is the predominant species. Subcluster Ba represents the link between wetter and drier localities because of the significant presence of *Pirata hygrophilus* (D cca 23 %). Subclusters Bb and Bc contain the communities with the highest dominance of the species *Ozyptila praticola* ($D > 31\%$) and with the lowest number of *Pirata hygrophilus* ($D < 10\%$).

Cluster C, with the communities of hard wood inundation forests, is distinguished from clusters A and B by the eudominant presence of the xerophilous species *Enoplognatha thoracica*, *Zelotes apricorum* and by the absence of *Pirata hygrophilus*.

The evaluated communities of the localities Gabčíkovo, Trstená na Ostrove and Vlčie hrdlo during the three years of research showed relative stability in composition and structure of the individual clusters. The most striking changes in the structure of the communities were found in the locality Bodíky, where a strong decrease of the hygrophilous species was found. This was manifested by the change in position of this locality in the dendrogram (shifting from subcluster Aa in 1991 to subclusters Bb and Bc in 1993 and 1992, respectively). The opposite tendency was registered in the locality Klúčovec, where a slight shifting towards the more wet habitat was observed during the research period. Small changes were also observed in the locality Dobrohošť, where the dominance of hemixerophilous species increased in comparison with the hygrophilous species during 1992–1993. These changes correspond with the soil moisture (Table 2) and Pišút's results. Pišút (1994), when monitoring of inundation forests, observed an obvious deterioration of the quality of woods and changes in species composition on the model areas of Dobrohošť and Bodíky. Similar changes were noticed in the quantity of Heteroptera in these model areas (Štepanovičová, 1994). This corresponds with our conclusions. Similar research was carried out by Krumpálová (1994). She studied the response of the arachnofauna to changes in the hydrological conditions in seven localities of inundation forest, near the Danube River during 1989–1991. She observed a decrease in the dominance of hygrophilous species *Pirata hygrophilus* during the research period and an increase of hemixerophilous species *Pardosa lugubris* and *Ozyptila praticola* which phenomenon she assumes to be connected with the changes in the level of ground water.

The research on the spider communities in inundation forests of the Danube River in Austria (Thaler & Steiner, 1987) and Germany (Bauchhenss, 1991) demonstrates the rather dry character of these habitats with a dominance of the hemixerophilous species *Pardosa lugubris* and *Ozyptila praticola*. Miller and Obrtel (1978) and Gajdoš (1994b) studied the spider communities in the inundation forests of the Morava River. These authors describe a similar composition of epigeic spider communities as near the Danube river, with a dominance of *Ozyptila praticola* in drier and *Pirata hygrophilus* in wet habitats.

Epigeic bog spiders for monitoring of habitat changes in Finland were used by Koponen (1979, 1980). No clear difference was found between the habitats in comparing the number of species, number of individuals and diversity. Changes in the

terrestrial spider fauna disturbed by human influence were studied in German raised bog after 22 years by Schikora (1994). On a general species level there were no clear clues ascertained about the consequences of biotope changes; 81% of the former species were found again. The spider *Antistea elegans* was discussed as a possible indicator species for monitoring hydrological changes in bog biotopes. According to our results occurrence of threatened and rare species and abundance of specimens of *Pirata hygrophilus*, *Ozyptila praticola* could be considered as the bioindicators for monitoring hydrological changes in inundation forests. The proportion in which these two species are present in soft wood inundation forests is a very good marker for moisture evaluation of the model areas.

ACKNOWLEDGEMENT. The author is indebted to Doc. RNDr. Oto Majzlan, CSc. for the use of spider material, to Dr. P. Degma, CSc. member of Department of Zoology, Comenius University for the dendrogram preparations and to Dr. P. J. van Helsdingen for language correction.

REFERENCES

- BAUCHHENS E. 1991: Die epigäische Spinnenfauna eines Auwaldgebietes der Donau im Landkreis Dillingen (Donau) Deutsch land. Bayern. *Arachnol. Mitt.* 2: 20–30.
- BÖCKH G. 1857: Über die Spinnen der Umgebung Presburgs. *Verh. Ver. Naturk. u. Hilfk. Pressburg* 2 (2): 72–86.
- BÖCKH G. 1862a: Über das periodische Erscheinen der Spinnen und das zweckmässigste Einsammeln derselben. *Corresp.-Sl. Ver. Naturk. Pressburg* 1 (1): 1–9.
- BÖCKH G. 1862b: Über das periodische Erscheinen der Spinnen und das zweckmässigste Einsammeln derselben. *Corresp.-Sl. Ver. Naturk. Pressburg* 1 (2): 45–60.
- CHYZER C. & KULCZYŃSKI 1891: *Araneae Hungariae*, I. Verlag Academiae scientiarum hungaricae, Budapest, 168 pp.
- CHYZER C. & KULCZYŃSKI 1894: *Araneae Hungariae*, II., pars prior. Verlag Academiae scientiarum hungaricae, Budapest, 151 pp.
- CHYZER C. & KULCZYŃSKI 1897: *Araneae Hungariae*, II., pars posterior. Verlag Academiae scientiarum hungaricae, Budapest, pp. 143–366.
- CHYZER C. & KULCZYŃSKI 1899: Arachnida. In Thalhammer J. (ed.): *Fauna Regni Hungariae, pars III.: Arthropoda*, Budapest, pp. 1–33.
- GAJDOŠ P. 1987: Pavúky (Araneae) Štátnej prírodnej rezervácie Ostrov Kopáč. [Spiders (Araneae) of State nature reservation Ostrov Kopáč.] *Ochrana prírody, Bratislava* 8: 255–273.
- GAJDOŠ P. 1994a: The epigeic spider communities of lowlands forests in the surrounding of the Morava river. *Ekológia (Bratislava)* 13, Supplement 1, pp.135–144.
- GAJDOŠ P. 1994b: Monitorovací výskum epigeických spoločenstiev pavúkov (Araneae) lužných lesov Dunaja. (Monitoring research of epigeic spider communities of the inundation forests of the Danube river). *Proceedings of the conference on Biota monitoring on the area of interest of the Gabčíkovo water dam*. Ústav zoológie a ekozoológie SAV, Bratislava, p.12 (in Slovak).
- GAJDOŠ P. & SVATONĚ J. 1993: The red list of spiders of Slovakia. *Boll. Acc. Gioenia Sci. Nat.* 26 (345), pp. 115–133.
- GAJDOŠ P., SVATONĚ J., ŽITŇANSKÁ O. & KRUMPÁLOVÁ Z. 1992: Spiders (Araneae) of the Danubian plain. *Entomologické problémy* 23: 28–60.
- JANIŠ S. 1991–1994: *Monitoring of soil moisture for Slov. Acad. Scien. in framework of monitoring of biota on the surroundings of Gabčíkovo dam*. Final reports from the years 1990–1993, Geofyzika, Bratislava.
- KOPONEN S. 1979: Differences of spider fauna in natural and man-made habitats in a raised bog. In: *The National Swedish Environment protection Board, Report PM II 51*: 104–108.
- KOPONEN S. 1980: Effects of drainage on the spider fauna (Araneae) of the ground-layer on mires. In Dindahl D. L. (ed.): *Soil Biology as Related to Land Use Practises*. Proceedings of the VII International soil Zoology Coll., Washington, pp. 441–443.

- KRUMPÁLOVÁ Z. 1994: Reakcia arachnofauny na zmeny hydrologických pomerov Dunajských lužných lesov. (The reaction of the arachnofauna to the hydrological changes of the inundation of the Danube River.) *Proceedings of the conference on Biota monitoring on the area of interest of the Gabčíkovo water dam*. Ústav zoológie a ekoszológie SAV, Bratislava, p. 22 (in Slovak).
- LISICKÝ M., KOZOVÁ M. & MIKLÓS L. 1988: *Návrh kontrolného a pozorovacieho systému vplyvu SVD Gabčíkovo-Nagymaros z hľadiska biológie a krajinnej ekológie. (Proposal of a controlling and monitoring system on the influence of the Gabčíkovo water dam from the point of biology and landscape ecology.)* Internal materials, Ústav zoológie a ekoszológie SAV, Bratislava, 37 pp.
- MARTIN D. 1991: Zur Autökologie der Spinnen (Arachnida: Araneae). I. Charakteristik der Habitatausstattung und Präferenzverhalten epigäischer Spinnenarten. *Arachnol. Mitt.* 1: 5–26.
- MILLER F. & OBRTEL R. 1975: Soil surface spiders in lowlands forest. *Acta Sc. Nat. Brno* 9 (4): 1–40.
- ORTVAY T. 1902: Pozsonyvármegye és a területén fekvő Pozsony, Na gyszombat, Bazin, Modor's Szentgyorgy városok állatvilága. (Animal life in the region of Bratislava, Tnava, Pezinok, Modra, Jur pri Bratislave). In Stampfel K. (ed.): *Allatraxji rész. Pozsony I. köt. Pozsony*, pp. 574–596.
- PIŠŤ P. 1994: Stav lužných lesov podľa údajov monitoringu bioty v druhom roku prevádzky vodného diela Gabčíkovo. (State of the inundation forests according to the monitoring of the biota in the second year of the functioning of the Gabčíkovo water dam.) *Proceedings of the conference on Biota monitoring on the area of interest of the Gabčíkovo water dam*. Ústav zoológie a ekoszológie SAV, Bratislava, p.31 (in Slovak).
- PLATNICK N. I. 1991: *Advances in Spider Taxonomy 1988-1991 (With Synonymies and Transfers 1940-1980)*. New York, 846 pp.
- PODANI J. 1980: Számítógépes programcsomag ökológiai, cönológiai és taxonómiai osztályozások végrehajtására. *Abstr. Bot.* 6: 1–158.
- SCHIKORA H. B. 1994: Changes in the terrestrial spider fauna (Arachnida: Araneae) of a North German raised bog disturbed by human influence. 1964–1965 and 1986–1987: A Comparison. *Memoirs of the Entomological Society of Canada* 169: 61–71.
- ŠTEPANOVIČOVÁ O. 1994: Heteroptera ako súčasť epigeónu lužných lesov záujmového územia VD Gabčíkovo. (Heteroptera as a part of epigeon of the inundation forests on the area of interest of the Gabčíkovo water dam.) *Proceedings of the conference on Biota monitoring on the area of interest of the Gabčíkovo water dam*. Ústav zoológie a ekoszológie SAV, Bratislava, p. 38 (in Slovak).
- THALER K., PINTER M. & STEINER 1984: Fallenfänge von Spinnen in den östlichen Donauauen (Stockerau, Niederösterreich). *Spixiana* 7: 97–103.
- THALER K. & STEINER H. M. 1989: Fallenfänge von Spinnen in abgedämmten Donau-Auen bei Wien (Österreich). *Sber. Österr. Akad. Wiss. Mathem.-naturwiss. Kl., Avt.* 1, 196: 323–339.