

Hendrik Segers

Laboratorium voor Ecologie der dieren, Zoögeografie en Natuurbehoud  
Ledeganckstraat 35, B-9000 Gent

Jean-Pierre Maelfait

Instituut voor Natuurbehoud van de Vlaamse Gemeenschap  
Kiewitdreef 3, B-3500 Hasselt

#### A COMPARATIVE STUDY OF THE SPIDER COMMUNITIES OF DIFFERENT WOODLAND HABITATS

##### Introduction

The Zoniën forest is the largest forest ( + 4300 ha.) on silty soils in Belgium. The forest is of special interest because of its history : it has never been under agricultural management. This has resulted in the preservation of the original pedological characteristics which have developed on the Weichelian loess deposits (Langohr & Cuykens, 1985). The forest was however heavily exploited during the wars of the 16<sup>th</sup> and 17<sup>th</sup> century. During the Austrian period (18<sup>th</sup> century) a start was made of a massive reforestation with beech. This resulted in a tradition of beechmonoculture and in the present day some 80% of the forest is made up of beech stands.

Because of water percolation the originally calcareous loess deposit was leached out upto a depth of 2 to 3 m. Probably due to periglacial outdrying, the soil is compacted betweed approximately 30 and 120 cm of depth (Fragipan) (Van Vliet & Langohr, 1981). Under the present-day climatic conditions this layer does not form a barrier for water percolation but, with the exception of a polygonal pattern of fissures existing in it, is impenetrable for root growth. Another problem for the older beech stands is compaction of the surface soil due to their accessibility for horse riding and heavy forest exploitation machines (De Meyer & Langohr, 1984). Both the subsurface and surface compaction are the main causes of important windthrow of older beeches. Liming has been tried out as an attempt to treat the surface compaction.

It is within the context of these problems that we have started an investigation of the soil fauna. We discuss hereafter some preliminary



Table 2 : Morphology of the soil in the sampled stands :

Depth of surface compaction and Fragipan horizon (in cm.)

Stand :	Surface compaction :			Subsurface compaction		
	Median	Median	Compaction			Fragipan
	upper limit	lower limit	None/Weak/Strong			depth
A	-	-	7	0	0	58
B	6	10	0	7	0	48
C	4.5	28	0	0	7	32
D	6.5	29	0	1	6	38
E	0	30.5	0	1	6	33
F	-	-	7	0	0	32
G	6	16	0	0	7	25
H	6	11	1	5	1	50
I	6	19	0	0	7	27
J	-	-	7	0	0	40
K	6	11	0	6	1	25
L	7	17	0	3	4	31

### Results and discussion

On the whole 140 species of spiders were caught. The faunistics of some of them was briefly discussed in Segers & Maelfait (in press.). In the present study we confine ourselves to a synecological analysis of the most abundant species, i.e. species caught in a number higher than the total number of traps used. (> 98). Based on the procentual abundance of each species over the fourteen sampling units, a detrended correspondance analysis (Hill, 1979) was performed. In table 3 we listed these species and the sampling units according to their scores obtained along the first axis of this analysis. Also mentioned are the procentual abundance of each species per sampling station (/year), the value of Levins' measure for habitat width (Levins, 1968), and the sum of the procentual abundancies per station.

We can discern three groups of species. A first group of species is formed by the species predominantly occurring in A (and B). A second group consists of species with no clear preference for one of some of the sampling units.

Most of these were also observed in large numbers during an investigation of a german beech wood by Dumpert & Platen (1985). The third group comprises species showing a preference for the stations at the other end of the variation axis, G and I.

A is an old stand of oak with a deep soil and without soil compaction. It has a rich spider fauna with a large relative abundance of some small Linyphiid spiders (group 1). For Monocephalus fuscipes it is known that it is bound to forests with fastly decomposing litter (mull humus) (Baert, 1981). The more direct reason for that habitat binding is presumably that these spiders feed on springtails, which only occur in large numbers in good decomposing litter.

Station G and I are old open stands of beech. Due to forest exploitation their soils have a pronounced surface compaction. Station I has been limed some 30 years ago. Although also quite rich the soil spider fauna of both these stands is very differently structured than that of A. The typical spiders (group 3) are well represented in the open stands of beech with a well developed herb vegetation : G, I, L and E. They are almost absent in the catches of station K which only differs from L by the absence of that herb layer. Both K and L are old stands with a heavy compaction of the upper soil due to horse riding. L has been limed some 15 years ago. A spider community (a soil fauna) associated with a better litter breakdown like we find in A has not resulted from that treatment. Poor spider communities are also found in the other monocultures of beech without any herb layer : D and H. The difference between D and C is interesting. Both are young plantations of former old beech stands with severe soil compaction (which is in both still present). In C however we have a richer spider taxocoenosis with a relatively large abundance of the species associated with a good litter breakdown. In spite of the litter composition and the absence of soil compaction we find in J and F poor spider communities and indeed also litter accumulation (mor humus conditions). The reason for this unexpected poverty is not yet clear. Both are however very closed stands with few light and rain penetrating to the litter layer.

## Conclusion

As a whole, the above cited results suggest that soil faunas associated with a good litter breakdown are only present under open, mixed stands. Liming does not seem to bring about such a fauna. This is therefore not a good solution to treat the compacted soils of the Zoniën forest. Establishment of mixed and not too dense stands is probably a better treatment. The presence of a larger quantity of dead wood might also help and that by its amelioration of the nitrogen fixation (Pritchett & Fisher, 1987).

## Acknowledgements

The authors wish to thank Dr. R. Langohr for his help during our investigations.

The first author acknowledges an I.W.O.N.L. grant (no. 860457).

## Litterature

- Baert, L. : Synoecologie van de spinnefauna van boshabitaten. Deel I. Kenmerkende spinsoorten van de verschillende bemonsterde boshabitaten (Ecology of Belgian spiders I). Bull. Ann. Soc. r. Belge Int., 117 : 45-68 (1981)
- De Meyer, H. & Langohr, R. : Het Zoniënbos, of de menselijke invloed op de natuur. WIELEWAAL, 50 : 357-365 (1984)
- Dumpert, K & Platen, R. : Zur Biologie eines Buchenwaldbodens 4. Die Spinnenfauna. Carolea, 42 : 75-106 (1985)
- Hill, M.O. : DECORANA -- A FORTRAN program for detrended correstandance analysis and reciprocal averaging. Ecology and Systematics, Cornell University, Ithaca, New York (1979)
- Langohr, R. & Cuyckens, G. : Een bos op lemen voeten : bodem en reliëf in het Zoniënbos. NATUURreservaten '85 : 132-139 (1985)
- Pritchett, W.L. & Fisher, R.F. : Properties and management of forest soils. John Wiley & sons, New York (Sec. ed.) 1987
- Segers, H. & Maelfait, J.P. : Faunistical observations on the spider fauna of the Zoniën forest (Belgium). Actes X. Coll. Eur. Arachnol. Rennes 1987 (in press)
- Van Vliet, B. & Langohr, R. : Correlation between Fragipans and permafrost with special reference to silty Weichelian deposits in Belgium and Northern France : CATENA 8 : 137-154 (1981)

Table 3 : Procentual abundance of the frequently caught species in the different sampled forest stands with indication of the sampling year (85'-86 : 1, 86-87 : 2)

stand :	A1	A2	B1	C1	E1	F1	F2	J2	D1	K2	H2	L2	G2	I2	habitat width (Levins)
species:	(* : Values under median)														
<i>Monocephalus fuscipes</i>	42.6	36.4	10.0	2.5	3.8	1.3			0.8	1.3					*0.173
<i>Tapinocyba insecta</i>	30.2	27.8	22.7	0.8	12.7	0.8	0.4						3.2	1.2	*0.209
<i>Lepthyphantes flavipes</i>	24.8	18.6	14.6	26.5	3.5	3.7	2.5	1.0	3.5	0.2		0.2	0.5	0.2	*0.329
<i>Waikenaeria cucullata</i>	14.2	12.9	12.3	2.6	21.4	24.6	7.1		0.6	0.6		1.3	1.3	0.6	*0.370
<i>Diplocephalus picinus</i>	19.4	29.0	7.1	2.8	1.7	4.2	5.3	7.8	0.7	0.3	2.7	2.9	7.7	8.3	0.541
<i>Pirata hygrophilus</i>	0.9	1.9			61.6	13.0	21.0		1.9						*0.126
<i>Coelotes inermis</i>	10.9	11.0	8.9	4.3	12.2	12.7	15.9	6.6	3.2		4.7				0.674
<i>Diplocephalus latifrons</i>	28.4	26.9	3.2	1.5	8.0			0.2	1.2	0.5	0.2	2.0	12.4	15.2	*0.347
<i>Saarietoa abnormis</i>	7.2	8.0	16.8	10.4	0.8	11.2	6.4	12.0	4.8		7.2	1.6	8.0	5.6	0.690
<i>Waikenaeria acuminata</i>	12.7	7.8	12.7	1.8	5.0	15.0	9.6	3.2	1.4	0.5	6.4	1.8	13.6	8.2	0.637
<i>Macrargus rufus</i>	2.7	6.8	5.6	0.8	8.9	8.3	16.9	10.0	3.6	8.2	11.2	6.2	4.9	5.7	0.869
<i>Coelotes terrestris</i>	5.7	3.8	4.5	3.4	11.3	8.9	9.4	5.7	4.0	17.0	4.6	13.4	4.5	3.7	0.742
<i>Centromerus aequalis</i>	5.0	7.9	4.5	11.8	9.9	9.9	5.0		11.8	2.0	11.8	2.0	13.3	5.0	0.720
<i>Lepthyphantes zimmermani</i>	11.2	2.9	6.5	6.0	2.4	5.9	5.2	3.5	10.4	12.7	4.5	10.8	10.1	7.5	0.850
<i>Waikenaeria corniculans</i>	10.1	3.4	2.5	2.7	11.2	6.7	4.3	5.4	3.4	1.6	27.1	2.9	10.1	8.5	0.567
<i>Histoona torpida</i>	1.7	0.5	4.0	7.0	6.3	12.3	8.0	2.5	4.9	8.1	16.8	10.8	14.5	2.3	0.718
<i>Micrargus herbigradus</i>	2.3	0.5	15.9	6.9	4.1	11.4	5.6	5.3	0.3	1.0	3.6		17.7	25.3	0.505
<i>Centromerus sylvaticus</i>	3.6	10.3	6.7	1.5	2.9	2.0	3.1	3.1	2.6	2.9	11.4	14.3	8.9	26.4	0.515
<i>Micrometa viaria</i>	1.5	1.4	4.1	3.2	7.8	6.9	7.4	6.2	3.2	9.0	8.0	2.9	18.8	19.5	0.644
<i>Waikenaeria atrothibialis</i>	2.7	1.8	0.9	1.8	9.8	3.6	0.9		1.8		7.1	0.9	45.5	23.2	*0.251
<i>Linyphia clathrata</i>	3.6	0.6	0.6	5.4	1.8	5.4	0.6			4.2	7.2	16.8	29.5	24.0	*0.376
<i>Trochosa terricola</i>	0.7	1.4	2.2	1.4	9.4	1.4	2.2	1.4	3.6	5.0	5.0	11.5	9.4	45.3	*0.317
<i>Pardosa lugubris</i>	3.8	1.8	0.9	4.7					1.8	0.4	2.5	1.6	3.4	14.0	63.3
<i>Diplostylella concolor</i>	0.7	1.1	1.1	1.8	0.4	0.7			0.7	1.8	0.7	47.8	43.3		*0.174
<i>Pachygnatha listeri</i>				2.3	0.8			0.8		0.8	0.8	2.3	12.7	79.6	*0.128
Total :	234	226	172	98	228	166	148	90	70	78	144	108	316	425	

Jocqué: Can you explain why *Pirata hygrophilus* is very common in a few stands whereas it was virtually absent in most others. Could the presence of open spaces in some of these stands provide an explanation?

Segers: So far, we cannot explain this phenomenon with certainty. We suppose that the presence of flacks of water could be the explanation, but these also occur where *P. hygrophilus* is rare. The stands where it is common indeed have open spaces, but other stands with open spaces are poor in *P. hygrophilus*.