

Temporal species serie of web-spiders (Arachnida: Araneae) as a result of pine tree bark-structure

par
Ulrich Simon

Summary: In 1987, one group of 40 years old pine-trees (*Pinus silvestris* L.) and another group of 120 years old pines were investigated with arboreal photo-electors and arboreal pitfall-traps, 2 meters above ground.

Species with a high number of individuals showed a close serie of peaks on the old stems. On the young pines, peaks appear at a different time and the number of species living there is lower.

This is discussed as a result of the well structured bark of old pines offering its inhabitants, conditions that even ground living spiders can use.

1. INTRODUCTION

There is a lot of research concerning the fauna of tree stems (e. g. FUNKE 1977b, FUNKE & SAMMER 1980, ALBERT 1976 and 1982). But only a few of them are concerned with functional aspects of the bark (NICOLAI 1986). For spiders habitat-structure is one of the most important abiotic factors: to fix the net, to be hidden from predators, because of microclimatical conditions, etc.

In this work the influences of *Pinus silvestris* L. age and of its bark structure on species composition and the activity of spiders during the year on the bark are to be shown.

2. MATERIAL AND METHODS

The investigation was carried out with arboreal photo-electors (after FUNKE 1971) and arboreal pitfall traps at two groups of pines each of three trees (*Pinus silvestris* L.). The sampling period did last from May 7, 1987 to December 10, 1987.

The trees of the first group were about 120 years old, the ones of the second about 40 years old. At a height of 2 meters, two, three or four of each traps were installed as a sleeve around the stem. The spiders were captured in saturated picric acid. Every week all traps were emptied, the animals were put into ethanol (70%) and determined. Spiders were determined using GRIMM 1985, LOCKET & MILLIDGE 1951/53, LOCKET, MILLIDGE & MERRET 1974, ROBERTS 1987, WIEHLE 1956.

3. RESULTS

One can see two seasons of spider-activity on pine trees stems, one in late spring and early summer, and one in autumn. Both of them last the same time, about 10 weeks (figure 1).

In the first season, the number of species with a proportion of more than 1% of all sampled adult spiders, is higher (15 species) than in the second season (4 species)(table 1). The series of species in both seasons are different comparing the old stems (figure 1a) with the young ones (figure 1b). The individual-rich species in spring and early summer show a very close serie of maxima in contrast to the species in autumn. As a result there is a peak every week for at least one of the species occuring on the stems of old pines in the first season. Since in autumn there are only a few species on the bark of pines the maxima of abundance are only every two weeks (figures 1a and b) or more. On the young pines the serie of species in spring is not similar and species have another row of appearance (figure 1b).

4. DISCUSSION

The results show the existence of limiting factors for the activity of spiders on pine-tree bark. Already known factors are temperature, climate and radiation (NICOLAÏ 1986). The two investigated stands possess only a few differences in temperature and atmospheric humidity. Global radiation is higher in the old stand (v. STÜLPNAGEL, pers. comm.).

Therefore the main factor for the observed difference in spider population must be the structure of the bark. Older pines possess a well structured bark with a lot of clefts and chinks. Spiders can use them to be hidden from predators (e. g. birds) or for the their microclimatical conditions (e. g. radiation may heat the bark and produce higher




Fig. 1 - The occurrence of adults spider on the stems of old pine trees (a) and young pines (b). Lines mean the appearance of the species during sampling period; triangles = maximum of species activity on the old pines - circles = maximum of species activity on the young pines.

Table I

Species found on bark of pines (more than 5 individuals), separated into the two seasons and the two aged tree-groups

name of the species	preferred habitat	old trees	young trees
appearing in spring			
<i>Theridion mystaceum</i>	trees	yes	yes
<i>Xysticus audax</i>	trees	yes	no
<i>Xysticus lanio</i>	tree/ground	yes	no
<i>Lepthyphantes flavipes</i>	ground	yes	yes
<i>Moebelia penicillata</i>	trees	no	yes
<i>Entelecara acuminata</i>	tree/ground	yes	no
<i>Meioneta innotabilis</i>	trees	yes	yes
<i>Textrix denticulata</i>	tree/ground	yes	no
<i>Coriarachne depressa</i>	trees	yes	yes
<i>Diaea dorsata</i>	tree/ground	yes	yes
<i>Theridion tinctum</i>	trees	yes	no
<i>Enoplognatha ovata</i>	ground	yes	no
<i>Philodromus aureolus</i>	trees	yes	no
<i>Clubiona brevipes</i>	trees	yes	no
<i>Trematocephalus cristatus</i>	trees	yes	no
appearing in autumn			
<i>Drapetisca socialis</i>	trees	yes	yes
<i>Steatoda bipunctata</i>	trees	yes	no
<i>Lepthyphantes minutus</i>	ground	yes	yes
<i>Centromerus sylvaticus</i>	ground	yes	yes

temperatures, but in the clefts humidity may be very high as well). This may lead to an addition of two positive factors for spiders!

Owing to this a high number of ground-living spiders (PLATEN 1984) are able to enter this habitat and add to the bark-living spiders (e. g. *Lepthyphantes flavipes*, *Centromerus sylvaticus*). The result is the described close serie of abundance's maxima. Further investigations will show until which level ground-living spiders can use bark as that supposed "supplementary habitat".

LITERATUR

- ALBERT, R. - (1976). Zusammensetzung und Vertikalverteilung der Spinnenfauna in Buchenwäldern des Solling. *Faun. ökol. Mitt.*, 5: 65-80.
- ALBERT, R. - (1982). Untersuchungen zur Struktur und Dynamik von Spinnengesellschaften verschiedener Vegetationstypen im Hoch-Solling. Hochschulslg. *Naturwiss. Biol.*, 16: 147 pp., *Diss. Freiburg i. Br.*
- FUNKE, W. - (1977b). Die Stammregion von Wäldern.-Lebensraum und Durchgangszone von Arthropoden. *Jber. naturw. Ver. Wuppertal*, 32: p. 45-50.
- FUNKE, W. & SAMMER G. - (1980). Stammaufbau und Stammanflug von Gliederfüßern in Laubwäldern (Arthropoda). *Entomologia Generalis*, 6 (2/4), p. 159-168.
- GRIMM, U. - (1985). Die Gnaphosidae Mitteleuropas (Arachnida, Araneae). *Abh. Naturwiss. Ver. Hamburg(NF)*, 26: 1-318.
- LOCKET, G.H. & MILLIDGE, A.F. - (1951/53). British Spiders. *Ray Society. London.* Vol. I + II, 310/449 pp.
- LOCKET, G.H., MILLIDGE, A.F. & MERRET, P. - (1974). British Spiders. *Ray Society. London*, Vol. III, 315 pp.
- NICOLAI, V. - (1986). The bark of trees: thermal properties, microclimate and fauna. *Oecologia*, 69: 148-160.
- PLATEN, R. - (1984). Ökologie, Faunistik und Gefährdungssituation der Spinnen (Araneae) und Weberknechte (Opiliones) in Berlin (West) mit dem Vorschlag einer Roten Liste. *Zool. Beitr.*, N. F. 28 (3): 445-487.
- ROBERTS, M.J. - (1987). The spiders of Great Britain and Ireland. *E. J. Brill. Leiden.* Vol. 2.
- WIEHLE, H. - (1956). Linyphiidae-Baldachinspinnen, *In: DAHL, F.*, "Die Tierwelt Deutschlands", 44: 1-337.

Ulrich SIMON

Institut für Biologie, sekr-fr1-1, Technische Universität Berlin
Franklinstrasse 28/29, D-1000 BERLIN 10